## II. Amendments to the Claims

Claims 1-6 (Cancelled).

7. (Currently Amended) The  $\underline{A}$  glycopeptide of claim 1 which is a compound of formula II:

$$H_3$$
C,  $H_3$ C,  $H_3$ C,  $H_3$ C,  $H_4$ C,  $H_4$ C,  $H_5$ 

wherein:

 $R^3$  is -OH;

R<sup>5</sup> is hydrogen;

R<sup>19</sup> is hydrogen;

 $R^{2\theta} \xrightarrow{is} R^{\pi} - Y - R^{\theta} - (Z)_{x}, R^{f}, -C(\Theta) R^{f}, \text{ or } -C(\Theta) - R^{\pi} - Y - R^{\theta} - (Z)_{x}; \text{ and }$ 

R<sup>3</sup>, Y, R<sup>b</sup>, Z, x, R<sup>f</sup>, R<sup>3</sup>, and R<sup>5</sup> have any of the values defined in claim 1;

R<sup>20</sup> is -CH<sub>2</sub>-CH(OH)CH(OH)CH<sub>2</sub>-Y-R<sup>b</sup>-(Z)<sub>x</sub> or -CH<sub>2</sub>-CH(OH)CH(OH)CH<sub>2</sub>-R<sup>17</sup>;

R<sup>17</sup> is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkynyl, substituted alkynyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, or heterocyclic;

Y is selected from the group consisting of oxygen, sulfur, -S-S-, -NR<sup>c</sup>-, -S(O)-, -SO<sub>2</sub>-, -NR<sup>c</sup>C(O)-, -OSO<sub>2</sub>-, -OC(O)-, -NR<sup>c</sup>SO<sub>2</sub>-, -C(O)NR<sup>c</sup>-, -C(O)O-, -SO<sub>2</sub>NR<sup>c</sup>-, -SO<sub>2</sub>O-, -P(O)(OR<sup>c</sup>)O-, -P(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -OP(O)(OR<sup>c</sup>)O-, -OP(O)(OR<sup>c</sup>)NR<sup>c</sup>-, -C(=O)- and -NR<sup>c</sup>SO<sub>2</sub>NR<sup>c</sup>-; each Z is independently selected from hydrogen, aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic;

R<sup>b</sup> is selected from the group consisting of a covalent bond, alkylene, substituted alkylene, alkynylene and substituted alkynylene, provided R<sup>b</sup> is not a covalent bond when Z is hydrogen;

each R° is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heterocyclic and -C(O)R<sup>d</sup>;

each R<sup>d</sup> is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkynyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic; and

x is 1 or 2;

or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof.

Claims 8-12 (Canceled).

- 13. (Currently Amended) A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 1 Claim 7.
- 14. (Currently Amended) The pharmaceutical composition of claim 13, which wherein the composition further comprises a cyclodextrin.

Claim 15 (Canceled).

- 16. (Original) A method of treating a mammal having a bacterial disease, the method comprising administering to the mammal a therapeutically effective amount of a glycopeptide of claim 7.
- 17. (Original) A method of treating a mammal having a bacterial disease, the method comprising administering to the mammal a therapeutically effective amount of a pharmaceutical composition of claim 13.
- 18. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is  $-CH_2-CH(OH)CH(OH)CH_2-R^{17}$  and  $R^{17}$  is alkyl.
- 19. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is  $-CH_2-CH(OH)CH(OH)CH_2-R^{17}$  and  $R^{17}$  is aryl.
- 20. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is  $-CH_2-CH(OH)CH(OH)CH_2-Y-R^b-(Z)_x$  and Y is -NH-.
- 21. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is  $-CH_2-CH(OH)CH(OH)CH_2-Y-R^b-(Z)_x$  and Y is oxygen.

22. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is  $-CH_2-CH(OH)CH(OH)CH_2-Y-R^b-(Z)_x$  and Y is sulfur.

23. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is  $-CH_2-CH(OH)CH(OH)CH_2-Y-R^b-(Z)_x$  and  $R^b$  is alkylene.

24. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is  $-CH_2$   $CH(OH)CH(OH)CH_2-Y-R^b-(Z)_x$  and Z is hydrogen.

25. (New) The glycopeptide of Claim 7, wherein R<sup>20</sup> is a group of the formula:

26. (New) The glycopeptide of Claim 7, wherein R<sup>20</sup> is a group of the formula:

27. (New) The glycopeptide of Claim 7, wherein R<sup>20</sup> is a group of the formula:

$$CH_2$$
  $CH_2$   $CH_2$   $CH_3$   $CH_3$   $CH_3$   $CH_3$ 

28. (New) The glycopeptide of Claim 7, wherein  $R^{20}$  is a group of the formula: